

Colonoscopy & EGD with Magcitrate

[Save Page as PDF](#)

MAGNESIUM CITRATE PREP

READ ALL INSTRUCTIONS A WEEK BEFORE YOUR PROCEDURE

If you start blood thinner or weight loss medications after scheduling your procedure or have questions, please call our office 303-788-8888 or visit www.southdenvergi.com.

7 DAYS PRIOR TO PROCEDURE

- Purchase 1 box of Simethicone/Gas-X 125 mg tablets (any color/flavor is OK). You will need a total of 4 tablets for your prep.
- STOP iron supplements one week before the colonoscopy as they can be constipating.
- We recommend you avoid foods with sunflower, pumpkin, chia, flax, hemp, sesame or other small seeds. If you do eat something containing seeds you do not need to worry, we can still complete your colonoscopy as scheduled.
- If you take BLOOD THINNERS we will obtain authorization from your prescribing doctor. We will call you with specific instructions. If we have not called you within 7 days of your procedure, please call our office immediately.
- If you take Warfarin, Coumadin or Jantoven, you must have your PT/INR checked the day before your procedure. We will give you specific instructions for this test.
- Take your heart and blood pressure medications as close to your regular schedule as possible. If you are

scheduled to take them within the 2 hours prior to the scheduled procedure time, please take them earlier than the 2 hours and with only a small amount of water.

- You may continue to take aspirin as prescribed.
- If you are scheduled to have anesthesia, the anesthesiologist may contact you to review your medical history, medications and allergies.
- If you are on medication for DIABETES, check with the doctor that prescribes it because you may need a dose change the day before or the day of your procedure.
- If you take **WEIGHT LOSS MEDICATIONS** such as GLP-1's or Phentermine you **MUST STOP** it for at least 7 days prior to the procedure.
- DO NOT STOP Contrave or Naltrexone.
- If you use inhalers please bring them with you. If you use a CPAP machine, please bring it with you only if you are scheduled at the hospital.
- Sedation: Conscious (goal is to keep you comfortable/you may or may not sleep) OR: Anesthesia/MAC (you will sleep) Both may leave you feeling groggy, dizzy, sleepy or nauseous. Vomiting may occur.

1 DAY BEFORE THE COLONOSCOPY

- Start on a CLEAR LIQUID diet when you get up and continue all day. Do not eat any solid foods.
- If you are scheduled after 12 noon, tomorrow, you may have a small serving, light breakfast today before 9AM – then start clear liquids.
- You must complete this prep even if you already have diarrhea.
- YOU MAY NOT HAVE anything red or purple (as they can stay in the GI tract and look like blood), milk, milk products, non-dairy creamers, juices with pulp, nectar or alcoholic beverages.
- YOU SHOULD DRINK 8 OUNCES OF CLEAR LIQUID EVERY HOUR

FROM THIS LIST:

- Clear broth (Chicken, vegetable or beef bouillon)
 - Broth or consommé
 - Tea
 - Coffee (No cream, milk or non-dairy creamer)
 - Kool-Aid (No red or purple)
 - Gatorade/PowerAde
 - White cranberry juice
 - Apple juice
 - Lemonade/ Limeade
 - Italian Ices / Popsicles
 - Jell-O
 - Hard candy
-
- If you are on medication for DIABETES follow your prescribing doctor's recommendations for these medications. You should drink clear liquids with sugar because you will not be eating. If you monitor your blood sugar continue to do this at your usual times. Check your blood sugar before you leave for the procedure. We ask that GLP-1 medications be held for 7 days. Please contact your prescribing provider's office for specific instructions.
 - Continue taking heart, blood pressure and time sensitive medication today and the day of the colonoscopy.
 - AT 6:00 P.M. THE NIGHT BEFORE your Colonoscopy, drink 1 $\frac{1}{2}$ (One and one half) bottles of Magnesium citrate.
 - IMPORTANT: You must drink TWO more 16-ounce containers of water over the next hour.
 - Chew 2 simethicone/gas x tablets.
 - You may continue to drink other clear liquids through the evening.

THE DAY OF THE COLONOSCOPY

- At 5 hours prior to procedure drink the remaining 1 $\frac{1}{2}$

(One and one half) bottles of Magnesium citrate. You will have 2 hours to finish drinking this dose and the additional water. Your stools should now be watery, clear/yellow. If your bottom is sore you may use a wet washcloth, A&D Ointment, Preparation H or a similar cream.

- **IMPORTANT:** You must drink (2) TWO more 16-ounce containers of water over the next hour.
- Chew 2 simethicone/gas-x 125 mg tablets.
- **AFTER 4 hours prior to procedure YOU MAY NOT DRINK ANY MORE LIQUIDS, INCLUDING WATER.**
- **ONLY EXCEPTION:** Take heart, blood pressure and anti-seizure medications with small sips of water at the time you normally would.